

When facing a difficult problem that you can't seem to solve on your own, turning to your peers for help can be a good way to get you thinking outside the box.



think antagonists

I often pose the following miniature case study to the students of my kinesiology class: A client comes into your office experiencing a decreased left rotation range of motion of the neck. I then ask each student, one at a time, to name a muscle that he or she would work to help this person, and it's fine for a student to name the same muscle that another student has already mentioned.

My reason for asking this question is to see how the students are connecting the information they learn about muscles in the science classroom with practical hands-on clinical application. Whether these students are early

It often gets stuck in the mind that the primary or only importance of a muscle is its mover actions. In reality, in the world of massage, what is most important is the muscle's antagonist function.

in their education or near the end, I receive mostly the same answers. The muscles they most often name are the right-sided sternocleidomastoid, trapezius, anterior scalene and cervical transversospinalis musculature (multifidus or rotatores), and the left-sided splenius capitis or cervicis, levator scapulae, or cervical erector spinae musculature (iliocostalis, longissimus, and spinalis). Their reasoning is that the right-sided muscles listed above are contralateral rotators, therefore left rotators. And the left-sided muscles listed above are ipsilateral rotators, therefore left rotators.

On one level, these responses make me quite happy because they

show that the students have successfully learned the science of the muscles. They have learned the names of many muscles of the neck, they have learned which ones are contralateral rotators and which ones are ipsilateral rotators, and they have learned to reason that right-sided contralateral rotator muscles do left rotation and left-sided ipsilateral rotators do left rotation. However, these answers point to a flaw when it comes to the critical reasoning necessary to apply this knowledge to clinical scenarios. While each student was able to successfully name a left rotator muscle of the neck, what they did not realize is that the cause of the client's problem is not the left rotator musculature!

Why? If a client cannot perform full left rotation range of motion of the neck, it is extremely unlikely that the reason is that left rotator muscles, which are the mover muscles of left rotation, are too weak. After all, the weight of the neck and head is not much to move; and this movement is not against gravity. Unless the client is very elderly or weakened by disease, these muscles should be able to create this joint action.

And, even if the cause is weak left rotator mover muscles, the clinical goal would be to strengthen them and that is not within the scope of license of most massage therapists. Strengthening weak musculature is the role of athletic and fitness trainers, Pilates and yoga instructors, physical therapists and chiropractic physicians.

How about if the left rotator mover muscles are tight instead of weak? That cannot be the solution either because tight muscles do not decrease their own joint motion. In fact, if the left rotator musculature were tight enough, the client might be stuck in a posture of left rotation, without the ability to rotate to the right.

Hopefully the answer is becoming clear. The cause of the client's inability to fully left rotate the neck is not the left rotator muscles; it is the right rotator muscles, which are the antagonists to the joint action of left rotation.¹ When a muscle is tight, it limits motion to the opposite side of the body from where it is located because

¹ It is easy to become too focused on the musculature of the body and look past other tissues that can cause restriction of joint motion. In reality, any taut soft tissue on the opposite side of the joint from the restricted motion (including joint capsules, ligaments, fascial planes, or even skin) could cause a decreased range of motion. Further, degenerative joint disease (osteoarthritic bone spurs), if advanced, can also interfere with normal joint motion.

To have access to the complete article, subscribe to Digital COMT.

CLICK HERE TO SUBSCRIBE!