

Palpation Assessment

by Joe Muscolino

"NEVER TREAT WITHOUT AN ASSESSMENT"

There is an old adage in the world of medicine - never treat without a diagnosis. A similar principle can be articulated in the world of massage therapy - never treat without an assessment. After all, if a physical assessment has not been made, how do you know what the appropriate or best treatment is for the client; what areas should be worked and for how long; what strokes should be used; how deep should the pressure be; should stretching be done or perhaps joint mobilisation?

If no assessment is made, the likely scenario is that the therapist will perform a cookbook massage, a canned routine that every client receives. Although a cookbook massage can be generally therapeutic, altering local blood circulation, affecting sympathetic/parasympathetic nervous system balance, and providing the warmth of touch, it will probably not be effective in treating a specific musculoskeletal condition. To accomplish this, clinical orthopaedic massage is needed, that is, massage that specifically addresses the musculoskeletal conditions of the client on the table at that moment. And this requires a competent physical assessment.

ASSESSMENT SKILLS

Competent orthopaedic assessment involves both the verbal history that covers the client's past and present status, and the physical assessment of the state of the client's tissues. Although massage therapists learn special assessment tests, such as Phalen's test for carpal tunnel syndrome or Wright's test for thoracic outlet pectoralis minor syndrome, by far the most important assessment skill that a massage therapist possesses is muscle palpation.

MUSCLE PALPATION: ASSESSING MUSCLE TISSUE

Muscle palpation assessment involves using touch to assess the health of a target muscle and its fascial tissues. Is the muscle tight or loose? Is it inflamed or painful to touch? Does it contain adhesions, trigger points, or taut and tender bands? Assessing the muscle to answer these questions can point you towards the most effective treatment for the client.

However, all of this is of limited value if we do not know the target musculature we are assessing. Although it is possible to perform massage stroke manipulation to musculature without knowing what muscles we are working, possessing this knowledge can make our work much more effective – identifying the muscle alerts us to the direction of the fibres which helps determine the preferred direction of our strokes. It also helps us to understand and be aware of precautions and contraindications in the area. Further, it is difficult to know how to effectively stretch a tight muscle if we do not know what it is. But, perhaps most importantly, if we do not know which musculature is unhealthy, it is not possible to determine why the client is experiencing the condition. Given that unhealthy musculature often results from overuse, without identifying the muscle we cannot reason backwards to determine what the client is doing in their daily life to create the problem. This also means we cannot give accurate and helpful postural and movement self-care advice for the client when they are at home, at work and engaged in sports or other hobbies.

MUSCLE PALPATION: LOCATING AND DISCERNING THE TARGET MUSCULATURE

The foundation of muscle palpation literacy lies in being able to accurately locate and discern the target musculature that is being worked. There is a science and an art to this aspect of muscle palpation. Following are the four major guidelines to follow when performing muscle palpation. Critically thinking through the application of these guidelines can greatly improve palpation skills.

Guideline 1 - Know the attachments of the target muscle

Knowing the attachments of the target muscle tells us where to place our palpating fingers. For example, if we know that the deltoid attaches from the clavicle and scapula to the deltoid tuberosity of the humerus, we also know that we need to place our palpating fingers on the proximal arm between those two attachment sites. If our target muscle is superficial and well developed, this may be all that we need to do to initially locate it. However, we still might not be sure of its exact borders. We might be able to locate the centre of the muscle belly but how will we know if we have strayed off the target muscle and onto adjacent musculature? And if the target muscle is deep and difficult to feel, simply placing our fingers between its attachments is clearly not enough. Therefore, we also need Guideline 2.

Guideline 2 - Know the actions of the target muscle

Knowing the actions of the target muscle provides us with the necessary information to ask the client to perform a joint action that will engage and contract that muscle. For example, we know that the deltoid abducts the arm at the glenohumeral joint so we can ask the client to perform this action and feel for the deltoid to palpably harden (Figure 1).



▲ **Figure 1:** Asking the client to abduct the arm at the glenohumeral joint engages and contracts the deltoid. From Muscolino JE: *The Muscle and Bone Palpation Manual, With Trigger Points, Referral Patterns, and Stretching*. St. Louis, 2009, Elsevier. Photo taken by Yanik Chauvin.

Continuing with this logic, we also know that the anterior fibres of the deltoid flex the arm at the glenohumeral joint so we can ask the client to flex the arm instead to locate the anterior deltoid.

Similarly, we can use extension of the arm for the posterior deltoid. If the deltoid hardens and the adjacent muscles and other soft tissues remain soft, then it is easier to discern its borders from adjacent soft tissues.

Guideline 3 - Choose the best action of the target muscle

Assuming that the target muscle has more than one action, it is not good enough to simply choose any of its actions to engage it. We need to choose the best action. This is where many errors in palpation protocols are made. The importance of this can be understood if we keep in mind that our goal is not just to have the target muscle contract but also to ensure that it is the only muscle that contracts. This way, it will be the only palpably hard structure amidst a sea of soft tissues. The goal of an isolated contraction of the target muscle is not always possible but most of the time it can be attained if the right action - the best action - is chosen.

To do this, we need to know the actions of the adjacent muscles so we can isolate an action for the target muscle. In other words, we need to find an action of the target muscle that is distinct from the actions of the adjacent musculature.

Pectoralis minor is a good example of this. The major action of the pectoralis minor is protraction of the scapula at the scapulocostal joint. But if we ask the client to protract the scapula, the overlying pectoralis major will also contract because it also protracts the scapula. So we need to find a distinct action of the pectoralis minor. The best action to isolate pectoralis minor is downward rotation of the scapula. The client can perform this by extending and adducting the arm at the glenohumeral joint (via scapulohumeral rhythm, extension and/or adduction of the arm requires downward rotation of the scapula) (Figure 2). Using this action, the pectoralis major remains relaxed and loose, and the contraction of the pectoralis minor can be felt through it.

Sometimes the best action varies depending on which part of the target muscle we are palpating. A good example of this is the fibularis longus (FL; formerly named the peroneus longus), located in the lateral leg between the extensor digitorum longus (EDL) which is anterior to it and the soleus which is posterior to it (Figure 3). The FL everts and plantarflexes the foot. So which action is best to choose? That depends on which part of the FL we are palpating. If we are palpating the anterior aspect of the FL, then we need to discern it from the adjacent EDL. In this case, if we ask the client to evert the foot, both muscles will engage (the EDL is also an evtor of the foot) and it will be difficult to discern the FL from the EDL. The better action to choose is plantarflexion of the foot. This will engage the FL, but the EDL will remain relaxed. However, if we are palpating the posterior aspect of the FL, adjacent to the soleus, plantarflexion of the foot is not the wise choice because the soleus also does this action. Here, eversion of the foot is the better choice because it will engage the FL but the soleus will remain relaxed and soft (the soleus is an invertor of the foot).




▲ **Figure 2:** Downward rotation of the scapula is the best action to choose to engage and palpate the pectoralis minor. From Muscolino JE: *The Muscle and Bone Palpation Manual, With Trigger Points, Referral Patterns, and Stretching*. St. Louis, 2009, Elsevier. Photo taken by Yanik Chauvin.

Guideline 4 - Add resistance

The point of Guidelines 2 and 3 is to make the target muscle engage and contract so that it hardens. However, if the joint action does not require a sufficiently strong enough contraction, the target muscle might not harden enough to be clearly palpable. This is especially true if the body part being moved is not very heavy or if the direction of motion is not against gravity. For this reason, guideline number 4 is to add resistance to the contraction. In Figure 1, the therapist can be seen adding resistance to the client's distal arm as she attempts to abduct it.

There are two important aspects to be aware of when adding resistance. First is that adding resistance is not a battle between the therapist and client to see who is stronger. The goal is to ask the client to try to move their body part and for the therapist to add just enough resistance to equal the client's contraction so that it is isometric.



If the resistance is not sufficient to clearly palpate the target muscle, then add more resistance by asking the client to press harder until you can feel its contraction. In some cases, it might be better to ask the client to press more gently. What is most important is that you are creative and try different degrees of resistance until you can clearly feel the contraction of the target muscle.

The second aspect of adding resistance during a palpation protocol is where you contact the client when adding the resistance. This is crucially important and is where more mistakes are made in palpation protocols than anywhere else. The rule to follow when contacting the client to add resistance is to never cross a joint that does not need to be crossed.

In the case of the deltoid, note that the therapist is adding resistance by contacting the client on the distal arm (See Figure 1). He is not crossing the elbow joint to contact the client's forearm or crossing the wrist joint to contact the client's hand.

The reason for this is clear if we keep in mind that the purpose of a palpation protocol is to feel and discern the target muscle from adjacent musculature. If the therapist crosses the elbow joint and contacts the client's forearm, the client will probably engage elbow joint musculature, whose bellies are in the arm next to the distal end of the deltoid (perhaps brachialis or triceps brachii, depending on whether the arm is medially or laterally rotated while abducting). This might cause the therapist's palpating fingers to move off the deltoid onto this other musculature, believing it is still deltoid because it is engaged and palpably hard.

Other examples are the pronator teres and brachioradialis of the forearm. Resistance to pronation of the forearm at the radioulnar joints for the pronator teres should be done by contacting the elbow and forearm, not pronating the forearm at the wrist joint.

To have access to the complete article, subscribe to Digital COMT.

CLICK HERE TO SUBSCRIBE!